

## A retrospective study of morphological pattern of anaemia in adults

Anuradha Shah<sup>1</sup>, Amit Shah<sup>2\*</sup>

<sup>1</sup>Associate Professor, Department of Pathology, Bidar Institute of Medical Sciences, Bidar, Karnataka, India

<sup>2</sup>Associate Professor, Maheswara Medical College, Isnapur, Pattanchuru, Telangana, India

Received: 04-09-2021 / Revised: 12-10-2021 / Accepted: 17-11-2021

### Abstract

**Introduction:** Anemia is a global public health problem affecting both developing and developed countries with major consequences for human health as well as social and economic development. Anemia is one of the common findings in patients irrespective of the presenting complaints and disease. Globally, anemia affects 1.62 billion people, which corresponds to 24.8% of the population. India is one of the countries with very high prevalence rates. NFHS-3 (National Family Health Survey) reveals the prevalence of anemia to be 70-80% in children, 70% in pregnant women and 24% in adult men[2]. **Materials and Methods:** A hospital based retrospective study was carried out at Department of Pathology, Bidar Institute of Medical Sciences, Bidar from 1st January 2019 to December 2020. 100 patients were evaluated for morphological pattern of anaemia based on red cell indices, peripheral smear, and leukocyte and platelet parameters. Anaemia was defined according to WHO criteria as haemoglobin concentration lower than 13 g/dL in men and 12 g/dL in women. Elderly was defined as a person 60 years and above according to WHO criteria. **Results:** The results of CBC and PBP showed that 29 (29%), 55 (55%), and 16 (16%) of the patients had normocytic normochromic, microcytic hypochromic and macrocytic pattern of anaemia respectively. The gender wise classification of the patterns revealed that 18 male patients and 14 female patients had normocytic normochromic patterns, while 52 female patients and 13 male patients had microcytic hypochromic patterns whereas 9 males patients and 7 cases of female patients showed macrocytic pattern [Table 2]. **Conclusion:** The microcytic hypochromic pattern of anaemia is highly frequent among this sample of patients while the macrocytic pattern of anaemia is the lowest. According to gender groups, microcytic hypochromic pattern of anaemia is more common among females; malnutrition, increase of blood loss due to pregnancy or menstruation, and lack of iron absorption are the main causes, while the normocytic normochromic anaemia is highly prevalent among males, which is mainly due to blood loss or chronic diseases.

**Keywords:** Anemia, WHO criteria, NFHS-3, CBC, PBP.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

### Introduction

Anemia is a global public health problem affecting both developing and developed countries with major consequences for human health as well as social and economic development. Anemia is one of the common findings in patients irrespective of the presenting complaints and disease. Globally, anemia affects 1.62 billion people, which corresponds to 24.8% of the population. India is one of the countries with very high prevalence rates. NFHS-3 (National Family Health Survey) reveals the prevalence of anemia to be 70-80% in children, 70% in pregnant women and 24% in adult men[2]. Symptomatology of anemia which may be specific or non-specific, depends on factors such as age of the individual, cardio-vascular reserve, the chronicity of the anemia, co-morbidities and others. Non-specific features of anemia are hypoxia related effects on organ systems especially heart, brain and muscles. They include easy fatigability, dizziness, fainting spells, malaise, palpitations and dizziness[3]. World Health Organisation (WHO) defines the lower limit of normal for hemoglobin concentration at sea level to be 12.0 g/dl in women and 13.0 g/dl in men. Normal hemoglobin distributions vary with age, sex, and physiological status, e.g., during pregnancy. Anemia is not a diagnosis itself, but it is a sign of underlying disease. Hence the evaluation of a patient with anemia is directed at elucidating the causes for the patient's reduced number and

morphological changes of red blood cells. An initial morphological classification of anemia with integration of red blood cell indices and morphological characteristics is probably most useful. The classification of anemia is also very important for the appropriate treatment of the underlying disease. Hence anemia is categorised by RBC size as microcytic, normocytic, or macrocytic.

Microcytic anemia associated with iron deficiency is the most prevalent micronutrient deficiency disease in the world affecting 2 billion people.<sup>4</sup> Anemia is diagnosed either by manual method ie peripheral blood smear examination or by automated hematology analyzer. The microscopic examination of a peripheral blood smear provides a wealth of information to the clinician. Various forms of anemia may actually be diagnosed from abnormal red cell morphology reported on a blood smear examination. The red cell morphology evaluation should include examination for deviation in size, shape, distribution, concentration of haemoglobin, colour and the appearance of inclusions[5].

### Materials and methods

A hospital based retrospective study was carried out at Department of Pathology, Bidar Institute of Medical Sciences, Bidar from 1st January 2019 to December 2020. 100 patients were evaluated for morphological pattern of anaemia based on red cell indices, peripheral smear, and leukocyte and platelet parameters. Anaemia was defined according to WHO criteria as haemoglobin concentration lower than 13 g/dL in men and 12 g/dL in women. Elderly was defined as a person 60 years and above according to WHO criteria. Anaemia was further graded into Mild, Moderate and Severe according to WHO criteria as under-

\*Correspondence

Dr. Amit Shah

Associate Professor, Maheswara Medical College, Isnapur, Pattanchuru, Telangana, India

E-mail: [docacs@gmail.com](mailto:docacs@gmail.com)

Mild	11-12.9 g/dL (Men) 11-11.9 g/dL (women)
Moderate	8-10.9 g/dL
Severe	< 8 g/dL

Microcytic anaemia was defined as MCV below 80 fl, normocytic as MCV between 80 and 100 fl and macrocytic anaemia by an MCV above 100 fl. Dimorphic anaemia was defined with a normal MCV but a raised RDW (normal – 11-15%). All were correlated with Peripheral smear examination.

#### Inclusion Criteria

Patients fulfilling the WHO criteria of anaemia haemoglobin of less than 13 gm/dL in males and less than 12 gm/dL in females.

#### Exclusion Criteria

Patient who's Hb is less than 13 gm% but has received blood transfusion in the last 3 months. Patients on treatment with iron, folic acid, B12 supplements.

#### Statistical Analysis

The data was collected and analysed using standard statistical chi – square test,  $p < 0.05$  statistically significant. Data was entered in Microsoft excel and analysis was done using SPSS version 22.

#### Results

The results of CBC and PBP showed that 29 (29%), 55 (55%), and 16 (16%) of the patients had normocytic normochromic, microcytic hypochromic and macrocytic pattern of anaemia respectively [Table 1]. The gender wise classification of the patterns revealed that 18 male patients and 14 female patients had normocytic normochromic patterns, while 52 female patients and 13 male patients had microcytic hypochromic patterns whereas 9 males patients and 7 cases of female patients showed macrocytic pattern [Table 2]. Other significant findings seen in this study among patients of anaemia are: pancytopenia 12 (12%), neutrophilia 10 (10%), PMN toxic granules 03 (03%), eosinophilia 01 (01%), lymphocytosis 10 (10%), and immature cells 03 (03%). Patients having pancytopenia and immature cells further advised for bone marrow examination to rule out and diagnosis of aleukemic leukemia, subleukaemic leukemia, leukemia and other causes of anaemia. 03 (03%) cases of anaemia in this study had already diagnosed leukemia.

**Table 1: Frequency of Morphological Patterns of Anaemia among Study Participants**

Morphological Patterns of Anaemia	Frequency (%)
Normocytic normochromic	29 (29%)
Microcytic hypochromic	55 (55%)
Macrocytic anaemia	16 (16%)
Total	100

**Table 2: Frequency of Morphological Patterns among Study Participants with regard to Gender Groups**

Morphological Patterns of Anaemia	Males	Females	Total
Normocytic normochromic	18	12	30
Microcytic hypochromic	13	42	55
Macrocytic anaemia	9	7	16

**Table 3: Other Significant Findings in Study Groups**

Pancytopenia	12 (12%)
Neutrophilia	10(10%)
Lymphocytosis	10 (10%)
PMN Toxic granules	03(03%)
immature cells	03(03%)
eosinophilia	01(01%)

#### Discussion

The most common morphological type of anaemia was microcytic hypochromic anaemia (55) [55%], which was in accordance with Joshi et al. (55.53%) and Ahmed et al. (56.6%). In a study done by M. Bukar et al 2009, Microcytic hypochromic anaemia was the commonest pattern with 64.9% followed by Dimorphic anaemia with 34.7% and Macrocytic anaemia with 0.4% [6].

Our study showed that the Microcytic hypochromic pattern of anaemia was more frequent than the macrocytic pattern. The gender wise classification of the patterns revealed a microcytic hypochromic pattern of anaemia was more common among females. Malnutrition, an increase of blood loss due to pregnancy or menstruation, and lack of iron absorption are the main causes [7,8]. In male patients, the frequency of the normocytic normochromic pattern of anaemia was high, which may be mainly due to blood loss or chronic diseases. In normocytic anaemia, the important issue is to identify the main cause and to prescribe the right treatment. The aetiologies of normocytic normochromic anaemia are nutritional, renal diseases (e.g., renal failure) and increased red cell destruction. In macrocytic anaemia, the main causes are a low level of Vitamin B12 and a low level of erythrocyte folate. The estimation of both patterns of anaemia will guide the proper diagnosis of macrocytic anaemia [9,10].

The most common cause of folate deficiency is either nutritional or malabsorption of this vitamin. In contrast, the deficiency of Vitamin B12 is almost resulting from malabsorption.

#### Conclusion

Iron deficiency anaemia is the most common type of anaemia seen mainly in women in the reproductive age group. In our study, adolescent women were found to have high prevalence of anaemia. Since anaemia is more prevalent among low socio-economic groups, higher diagnostic tests lead to financial burden to the patients. The study of basic blood parameters and peripheral smear examination are less expensive and mandatory in the work up of anaemia.

Morphological patterns of anaemia reflect the underlying aetiology, the study of which would ensure benefits in the early detection and appropriate treatment. Iron deficiency anaemia being a nutritional anaemia, can be prevented by improving nutritional status, creating awareness, and educating people.

#### References

1. Swaroop Raj BV, Sunita BS, Vanmalinitewari, Divya C. Is Peripheral Blood Smear an Obsolete Art? Int J Rec Scie Res 2015; 6(3): 3059-62.
2. Adewoyin AS, Ogbenna AA. Morphologic Evaluation of Anemia – I. Biol Med Aligarh 2016; 8(6):322.
3. Ejaz F, Ahmed M, Hassan NA. Comparative study in the diagnosis of anemia by Sysmex KX-21N hematology analyzer with peripheral blood smear. Int J Endor Health Sci Res 2013; 1(2):89-92.
4. Jones KW. Evaluation of cell morphology and introduction to platelet and white blood cell morphology. In: Harmening D (ed) Clinical Hematology and fundamentals of hemostasis, Philadelphia, 2009: 93-116.
5. Jadhav MV, Agarwal SA, Kadgi NV, Khedkar SS, Kulkarni KK, Kudrimoti JK. Utility of automated RBC parameters in evaluation of anemia. Int J Healthcare Biomed Res 2015; 3(3):170-81.
6. Singhal S, Verma N, Rathi M, Singh N, Singh P, Sharma SP, et al. Can peripheral blood smear examination be totally replaced by automated hematology analyser – with special reference to anemia? Int J Res Med Sci 2016; 4(10): 4563-66.
7. Sandhya V, Rashmi GS. Correlation of peripheral smear with RBC indices and RBC histograms in the diagnosis of anemia. Indian Journal of Pathology and Oncology 2017; 4(2):242-6.
8. Mishra P, Ahluwalia SK, Garg PK, Kar R, Panda GK. The Prevalence of Anemia among Reproductive Age Group (15-45yrs) Women in a PHC of Rural Field Practice Area of MM Medical College, Ambala. India J Women's Health Care 2012; 1:3.
9. Mukaya JE, Ddunga H, Ssali F, Shea T, Crowther MA. Prevalence and morphological types of anaemia and hookworm infestation in the medical Emergency ward, Mulago Hospital, Uganda. S Afr Med J 2009; 99:881-886.
10. Chandurkar M, Pendalya S. Clinical profile of patients presenting with anemia in a tertiary care rural hospital of Western Maharashtra. Int J Clin Biomed Res 2017; 3(4):74-77.

**Conflict of Interest: Nil Source of support: Nil**