

## A prospective study of knowledge, awareness, and practice of different nerve blocks for mandibular molar root canal treatment

Sudhakar Rao Malapati<sup>1\*</sup>, K.V.Kishore<sup>2</sup>, Kodali Sri Supraja<sup>3</sup>

<sup>1\*</sup>Oral and maxillofacial Surgeon, Santhiram Medical College and Hospital, Nandyal, India

<sup>2</sup>Craniofacial Surgeon, Santhiram Medical College and Hospital, Nandyal, India

<sup>3</sup> Oral and Maxillofacial Pathology, Dr Sudhakars Dental World, Nandyala, Andhra Pradesh, India.

Received: 08-12-2020 / Revised: 14-01-2021 / Accepted: 26-01-2021

### Abstract

**Introduction:** Local anaesthetics are the most widely and commonly used drugs in dentistry. Pain and dentistry are often synonymous in the minds of patients, particularly those with poor dentition due to various extractions, periodontal disease requiring surgery or symptomatic teeth requiring endodontic therapy. Dental practitioners, before beginning the procedure, identify a good anaesthetic agent and technique to focus solely on the operative procedures without distractions from pain-induced patient movements. **Materials and Methods:** study population that was included as part of the study included practitioners in the field of endodontic and postgraduates in the field of endodontic as these are the participants who are primarily involved in administering Local Anaesthesia during a root canal treatment on an everyday basis. The study was circulated in the form of an online survey collecting responses from the participants. **Results:** 50 % of the respondents administered a classical inferior alveolar nerve block involving the IAN, lingual and long buccal nerve. 26 % of the respondents anesthetized the inferior alveolar nerve and lingual nerve and 17 % anesthetized the inferior alveolar nerve and long buccal nerve. **Conclusion:** Anaesthesia of IAN alone should suffice during RCT of the mandibular molars. However, there is no adequate information and knowledge regarding the technique and the amount of local anaesthetic to be delivered for root canal therapy for mandibular molars thus inflicting further pain to the patient.

**Key Words:** Local anaesthetics, Pain, Dentistry, IAN.

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

### Introduction

Local anaesthetics are the most widely and commonly used drugs in dentistry. Pain and dentistry are often synonymous in the minds of patients, particularly those with poor dentition due to various extractions, periodontal disease requiring surgery or symptomatic teeth requiring endodontic therapy[1]. Dental practitioners, before beginning the procedure, identify a good anaesthetic agent and technique to focus solely on the operative procedures without distractions from pain-induced patient movements[2]. Research has shown that the major fear during a dental procedure is the fear of pain associated with intraoral administration of local anesthetics, which is the most common method for blocking pain during dental procedures. This is considered aversive due to the pain associated with the injection and the perceived threat of needle puncture prior to the injection[3]. Another survey finding was that those individuals who reported as highly fearful of dental procedures were worried about receiving oral injections and demonstrated an association between high dental anxieties and missed or / and delayed appointments. There is no doubt that the most widely used method for pain management during dental, and particularly endodontic, procedures is to administer intraoral local anaesthesia. Pain management approaches during root canal therapy can be based on one or a combination of such mechanisms. Blocking the nociceptive impulses during root canal treatment is performed with the administration of local anaesthesia[4]. Pain management during and after root canal treatment is a major challenge for dental practitioners [5,6]. Various studies of investigations have been

performed to overcome pain during and following root canal therapy[7]. Patient care should always put as the first priority during any clinical procedure. However the correct anaesthetic protocol to be followed for a root canal treatment has not been adequately documented, this further leads to various practices that lead to issues regarding the quality of life of the patient. This is a first of its kind study where the anaesthetic requirement for a root canal treatment is being assessed. One of the most common procedures that is carried out in the field of Endodontics is root canal treatment and hence appropriate administration of anaesthesia and reducing pain is of utmost importance in the field. The aim of the study is to assess the knowledge awareness and practice of different nerve blocks administered during Mandibular molar root canal treatment.

#### Materials and methods

The questionnaire was sent to 80 participants and all responded. In the present KAP survey, the study population that was included as part of the study included practitioners in the field of endodontic and postgraduates in the field of endodontic as these are the participants who are primarily involved in administering Local Anaesthesia during a root canal treatment on an everyday basis. The study was circulated in the form of an online survey collecting responses from the participants. This is a first of its kind study and thus the questionnaire was unique for the same.

Various questions that were included as part of this study were:

1. Employment Status as an Endodontist.
2. What nerve block do they give for the root canal treatment of a Mandibular Molar?
3. How much ml of LA do they use for a lower molar RCT.?
4. What type of LA do they use?
5. What Branches do they anaesthetise for a lower molar RCT.?

\*Correspondence

Dr. Sudhakar Rao Malapati


Oral and maxillofacial Surgeon, Santhiram Medical College and Hospital, Nandyal, India

E-mail: [sudhakarraomalapati@yahoo.com](mailto:sudhakarraomalapati@yahoo.com)


- 6. In 10 cases of lower molar RCTs, how many cases do they repeat the block.
  - 7. How many cases do they give an Intra pulpal dose of LA as well in about 10 cases?
  - 8. How effective do they think their block is?
  - 9. Do they accompany their block with an infiltration?
- In which state of the treatment does the patient experience maximum pain?

**Results**

A total of 80 responses was received. The responses were then verified and the results were tabulated. The results were plotted in the forms of graphs and then analysed. Out of the study population, 39 % of the participants were practitioners and 61 % of the participants were post graduate students in the field of endodontics. (Fig. 1)




**Fig 1: Employment Status as an Endodontist**



**Fig 2: Type of Local Anaesthesia Used**

A question was posed as to what nerve block, should they give for a lower molar root canal treatment and the entire study population (100 %) had opted for classical nerve block, none of the study participants opted for Gow Gates or Vazirani technique. The most commonly used type of LA is incorporated along with adrenaline through the responses received as part of the present study as 100 % of the participants opted for using lignocaine along with adrenaline. (Fig. 2) The question which was later posed to the study population involved the amount of local anaesthetic they would use for a lower molar for a root canal treatment. (Fig. 3) shows that 65.8 % of the population use about 2 – 3 ml, 31.6 % of the population use 1 – 2 ml and the

remaining 2.6 % use 3 – 4 ml of local anaesthetic agent. The responses that were established through this question is of high importance. (Fig. 4) 50 % of the population anaesthetise the inferior alveolar nerve along with the long buccal and the lingual nerve. On Probing on the use of an infiltration during the root canal treatment (Fig. 5), 50 % of the population admitted to administering it along with a nerve block. The participants were then asked in about 10 cases how many they would administer an Intra pulpal dose of local anaesthetic in. (Fig. 6) Majority of the population (34.2 %) have opted that for about 2 cases they would administer an Intra pulpal dose of LA as well.



**Fig 3: Quantity of LA Used for Lower Molar RCT**




Fig 4: Branches of IAN That are Anaesthetized




Fig 5: Whether Nerve Block is Accompanied with Infiltration

## Discussion

Local anaesthesia administration is a crucial aspect that is to be considered in patient-care to deliver painless treatment procedures. This is a first of its kind study. In the present study it is observed that for a lower molar root canal treatment practitioners relied more on classical inferior alveolar nerve block than other techniques. This can be attributed to the fact that the method is the most widely taught method of inferior alveolar nerve block administration and the ease of the same.<sup>8</sup> Although Gow Gates uses a more geometric and mathematical approach based on planes and provides good success rates the same has not been incorporated into clinical practice<sup>[9]</sup>. All participants voted that they administer local anaesthetic in combination with adrenaline. The use of adrenaline prolongs the duration and depth of anaesthesia as well as helps to have a better onset of action of the same<sup>[10]</sup> and hence it can be considered beneficial that the present study population prefer that over the alternative without adrenaline. But it has to be taken into account that when lignocaine is used along with adrenaline there is a significant change in the blood glucose level that is observed and adequate precautions have to be taken regarding the same in patients with systemic conditions such as diabetes. There could be a number of reasons for failure of pulpal anaesthesia using Inferior alveolar nerve block such as altered membrane excitability of peripheral nociceptors or due to Tetrodotoxin resistant channels which are basically a class of sodium channels that are resistant to the action of the local anaesthetic agent, a related factor is the increased expression of sodium channels in pulp diagnosed with irreversible pulpitis, psychological factors or due to the most common factor which is hot tooth. There are various strategies to tackle a hot tooth situation and the administration of Intra pulpal anaesthesia is definitely one of them and hence the candidates are adequately informed regarding the

same. From the above discussion it is very evident that unnecessary pain is being inflicted on the patient which requires prompt clinical intervention. Further studies are to be performed to standardize the type of nerve block that is to be administered for the particular procedure.

## Conclusion

Root canal treatment is the most commonly performed endodontic procedure and the administration of proper LA and creating a painless experience for the patient is very important. The inferior alveolar nerve block alone is sufficient for a lower molar root canal treatment but the knowledge regarding the amount of LA and type of block to be administered is not documented. This is currently leading to unnecessary pain being inflicted on the patient.

## References

1. Hamedy R, Shakiba B, Fayazi S, et al. Patient-centered endodontic outcomes: a narrative review. *Iran Endod J* 2013;8(4):197-204.
2. Eghbal MJ, Torabzadeh H. Review of high level endodontic research in pubmed index journals from Iran. *Iran Endod J* 2012;7(3):109-14.
3. Khalil H. A basic review on the inferior alveolar nerve block techniques. *Anesth Essays Res* 2014;8(1):3-8.
4. Kafalias MC, Gow-Gates GA, Saliba GJ. The Gow-Gates technique for mandibular block anesthesia. A discussion and a mathematical analysis. *Anesth Prog* 1987;34(4):142-9.

- 
5. Saxena A, Srivastava S, Agrawal A, et al. Comparison and clinical efficacy of local anesthetic solution xylocaine with and without adrenaline [1: 200000] in dental extraction. Saint's Int Dent J 2015;1(2):96-100.
  6. Ketabi M, Shamami MS, Alaie M, et al. Influence of local anesthetics with or without epinephrine 1/80000 on blood pressure and heart rate: a randomized double-blind experimental clinical trial. Dent Res J (Isfahan) 2012;9(4):437-40.
  7. Reader AL, Nusstein J. Local anesthesia for endodontic pain. Endod Topics 2002;3(1):14-30.
  8. von Arx T, Nakashima MJ, Lozanoff S. The face - a musculoskeletal perspective. A literature review. Swiss Dent J 2018;128(9):678-88.
  9. Jääskeläinen SK, Forssell H, Tenovu O, et al. Difficult diagnosis of facial pain: a case report and mini-review. Scand J Pain 2010;1(4):179-83.
  10. Drum M, Reader A, Beck M. Long buccal nerve block injection pain in patients with irreversible pulpitis. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2011;112(1):e51-4.

**Conflict of Interest:** Nil

**Source of support:** Nil