

## A retrospective study of correlation of risk factors with severity of diabetic retinopathy

Battula Yallamanda Babu Rao<sup>1\*</sup>, Rajalingam Vairagyam<sup>2</sup><sup>1</sup>Associate Professor, Department of Ophthalmology, Maheshwara Medical College and Hospital, Chitkul (V), Near Isnapur X Roads, Patanchervu(M), Sangareddy Dist, Telangana, India<sup>2</sup>Professor and HOD, Department of Ophthalmology, Superintendent, Sarojini Devi Eye Hospital, Humayun Nagar, Mehdiapatnam, Hyderabad, India

Received: 05-12-2020 / Revised: 10-01-2021 / Accepted: 01-02-2021

**Abstract**

**Introduction:** Diabetes mellitus is increasing in our society, so are the complications of diabetes mellitus. Diabetic retinopathy is a common cause of preventable blindness. We have taken a total of 1053 diabetic patients in our tertiary centre and studied 302 patients of Diabetic retinopathy diagnosed in the Department of Ophthalmology as a cross-sectional institution-based descriptive study from April 2016 to May 2018 in our tertiary care centre. **Materials and Methods:** We studied 302 patients of diabetic retinopathy and graded them into five grades of retinopathy. The severity of retinopathy is correlated with age, sex, duration of diabetes mellitus, systolic and diastolic blood pressure, BMI, albuminuria, smoking status and alcohol consumption. **Results:** Among the 302 diabetic retinopathy patients, there is a male predominance among diabetics. Females outnumbered males in the diabetic retinopathy group. Female preponderance in DR is not significant. There is an age-related increase in diabetic retinopathy, but the values are not statistically significant. 60% of our patients of DR are agricultural labourers in our study. Average age of diabetic retinopathy patients in our study is around 54. Duration of diabetes increased the prevalence of DR, but the values are not significant in our study. Systolic blood pressure, albuminuria and smoking history correlated with severity of DR (p-value < 0.05). BMI, alcoholism and diastolic blood pressure did not correlate with severity of diabetic retinopathy. **Conclusion:** There is a statistically significant correlation between severity of Diabetic retinopathy and systolic blood pressure, albuminuria and smoking. Other factors like female preponderance, higher BMI, diastolic blood pressure and alcohol intake were high among patients of diabetic retinopathy, but the values did not show statistically significant correlation with the severity of diabetic retinopathy.

**Keywords:** Diabetes mellitus, BMI, DR, diastolic blood pressure, albuminuria.

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

**Introduction**

Diabetes mellitus is the most common endocrine disorder. A report by International Diabetes Federation (IDF) on the global burden of Diabetes highlighted that-

1. 382 million people have diabetes in 2013; by 2035 this will have risen to 592 million.
2. The greatest number of people with diabetes is between 40-59 years of age and 80% diabetics are in low and middle income countries.
3. 175 million people with diabetes are undiagnosed.
4. More than 79000 children developed Type 1 diabetes in 2013.

Diabetic retinopathy has been classified by various methods, but the most commonly accepted [1-3]

Classification is according to International Clinical Diabetic Retinopathy Scale which classifies diabetes retinopathy into Non Proliferative (NPDR) and Proliferative Diabetic Retinopathy (PDR).

It is not possible to define which diabetic individual will present with retinopathy. However it is possible to determine the risk factors related to development of retinopathy, such as duration of disease, poor metabolic control, dyslipidemia, hypertension, alcoholism, associated renal disease, anaemia, hypomagnesaemia and smoking. Among these, duration of diabetes is proposed to be the most

significant risk factor.

**Materials and methods**

**Study Design:** It is a cross-sectional institution-based descriptive study from April 2016 to October 2017 in a tertiary care centre.

**Inclusion Criteria**

1. All the patients diagnosed as Diabetic Retinopathy above 20 years of age.
2. Patients of both sexes are taken into the study.

**Exclusion Criteria**

Non-cooperative patients and patients with Diabetic retinopathy who are seriously ill with other complications.

**Methodology**

Among a total of 335 diabetic retinopathy patients, 33 patients were excluded because of non-cooperation or seriousness of associated illness. 302 patients of Diabetic retinopathy were included in the study. All the 302 patients with diabetic retinopathy were subjected to dilated funduscopy with 90+ D lens direct ophthalmoscopy and were graded as follows-

- > Microaneurysm(s) only- Grade I.
- > Mild NPDR- Grade II.
- > Moderate NPDR- Grade III.
- > Severe NPDR- Grade IV.
- > Proliferative DR- Grade V[5-7]

Prevalence of Diabetic retinopathy, age, sex, occupation, duration of diabetes, BMI, systolic and diastolic blood pressure, smoking history and alcoholic history of these patients were studied. Routine blood and urine examination was done. Presence of albuminuria was noted.

**\*Correspondence**

**Dr. Battula Yallamanda Babu Rao**

Associate Professor, Department of Ophthalmology, Maheshwara Medical College and Hospital, Chitkul (V), Near Isnapur X Roads, Patanchervu(M), Sangareddy Dist, Telangana, India

E-mail: [raoy9@hotmail.com](mailto:raoy9@hotmail.com)

All the parameters were correlated with the severity of Diabetic retinopathy.

**Statistical Analysis**

Descriptive statistics were used to present the findings. Chi-square test of association was used to study the association between severity of retinopathy and different variables under study. Pearson’s Chi-square test is used as statistical hypothesis test and Chi-square

calculator for a contingency table that has up to five rows and columns is used for calculation. IBM Corp. Released 2016. IBM SPSS Statistics for Windows, Version 24.0. Armonk, NY: IBM Corp. is used as the statistical tool[8-10]

**Results**

Male Diabetics 557 and Female Diabetics 496. Ratio is 52.89%: 47.11%. Average age of diabetic retinopathy patients is 54.44

**Table 1: Age Distribution of Diabetic Retinopathy Patients**

Grade of Retinopathy	20-30 Years	30-40 Years	40-50 Years	50-60 Years	> 60 Years	Total
Grade I	02	03	22	39	18	84
Grade II	04	06	26	49	19	104
Grade III	01	12	21	42	06	82
Grade IV	02	03	02	04	05	16
Grade V	01	01	02	09	03	16
Total	<b>10</b>	<b>25</b>	<b>73</b>	<b>143</b>	<b>50</b>	<b>302</b>
Grade I	02	03	22	39	18	84

**Table 2: Sex Distribution of patients of Diabetic Retinopathy**

Grade of Retinopathy	Males	Females	Total	Percentage
Grade I	36	48	84	27.81%
Grade II	49	55	104	34.44%
Grade III	43	39	82	27.15%
Grade IV	08	08	16	5.29%
Grade V	08	08	16	5.29%
Total	<b>144</b>	<b>158</b>	<b>302</b>	<b>100</b>
Percentage	<b>47.68%</b>	<b>52.32%</b>	<b>302</b>	<b>100</b>

**Table 3: Professional Occupational Analysis of Diabetic Retinopathy Patients**

Profession	No. of Patients	Percentage
Agricultural Workers	181	59.93%
Skilled Labourers	076	25.16%
Others	045	14.90%

**Table 4: Duration of DM and Grade of Retinopathy**

Grade of DR	0-5 Years	5-10 Years	10-15 Years	>15 Years	Total
Grade I	11	13	28	32	84
Grade II	09	21	39	34	104
Grade III	16	16	28	22	82
Grade IV	02	02	09	03	16
Grade V	03	04	07	02	16

**Table 5: Systolic Blood Pressure and Diabetic Retinopathy**

Grade of DR	SBP 100-130 mmHg	SBP 130-160 mmHg	SBP 160 to 190 mmHg	SBP > 190 mmHg	Total
Grade I	22	16	36	10	84
Grade II	17	29	42	15	104
Grade III	14	32	25	10	82
Grade IV	02	05	04	05	16
Grade V	02	05	03	06	16

**Table 6: Diastolic Blood Pressure and Diabetic Retinopathy**

Grade of DR	DBP < 90 mmHg	90 to 100 mmHg	100 to 110 mmHg	> 110 mmHg	Total
Grade I	11	27	32	14	84
Grade II	21	21	39	22	104
Grade III	19	21	29	13	82
Grade IV	02	04	04	06	16
Grade V	02	04	04	06	16

**Table 7: Diabetic Retinopathy and BMI**

Grade of Retinopathy	BMI < 20	20-25	25-30	> 30	Total
Grade I	06	15	31	32	84
Grade II	18	13	34	38	104
Grade III	22	11	27	22	82
Grade IV	03	02	04	07	16
Grade V	04	02	05	05	16

**Table 8: Albuminuria and Diabetic Retinopathy**

Retinopathy	Patients with Albuminuria= 264	Patients with No Albuminuria= 38	Total 302
Grade I	76	08	84
Grade II	93	11	104
Grade III	74	08	82
Grade IV	09	07	16
Grade V	12	04	16

**Table 9: Smoking and Diabetic Retinopathy**

Retinopathy	Non-Smokers 92	Ex-Smokers = 154	Current Smokers = 56	Total 302
Grade I	32	36	16	84
Grade II	18	63	23	104
Grade III	31	42	09	82
Grade IV	05	06	05	16
Grade V	06	07	03	16

**Table 10: Alcohol Consumption and Diabetic Retinopathy**

Retinopathy	Alcohol Consumption Not Present = 273	Alcohol Consumption Present = 29	Total =302
Grade I	77	07	84
Grade II	92	12	104
Grade III	77	05	82
Grade IV	13	03	16
Grade V	14	02	16

31.81% of the diabetes mellitus patients were found to have Diabetic retinopathy. Average age of DR patients is 54 years. Predominant number of patients of DR are females. Female: Male Ratio is 52.32:47.68, though there is male preponderance among diabetic patients 52.89: 47.11. But the value is not statistically significant. Diabetic retinopathy occurred in predominantly above 40 years' age group. There is a statistically significant increase in grades of retinopathy with increased age. 60% of the patients in our study are agriculture workers and 25% skilled laborers. In our study, prevalence of DR increased with increase in duration of Diabetes mellitus. But the value is not statistically significant. There is a statistically significant increase in prevalence of DR with increase in systolic blood pressure. There is an increase in prevalence of DR with increase in Diastolic blood pressure, but the value is not statistically significant. 195 cases out of a total of 302 cases belong to group of 25 and above BMI. But the value in our study is not statistically significant. Albuminuria in our study statistically correlated with the severity of Diabetic retinopathy. Smoking correlated with severity of Diabetic retinopathy. History of alcoholism did not correlate with severity of Diabetic Retinopathy

#### Discussion

Though the number of diabetics were more among the men, retinopathy is predominant among women in our study. A Pakistani study also showed female preponderance. The study showed a

prevalence of 23.9% of DR, which is similar to our study. 17 Female predominance in our study is not statistically significant. A large Japanese study also showed female predominance. 29.6% of their diabetic patients had retinopathy. Another Indian study showed that duration of Diabetes and Diastolic Blood pressure showed a positive association with diabetic retinopathy with male predominance. The prevalence of DR in the Chennai Urban Rural Epidemiology (CURES) Eye Study in south India was 17.6 percent. CURES Eye study showed that the major systemic risk factors for onset and progression of DR are duration of diabetes, degree of glycaemic control and hyperlipidaemia. Hypertension did not play a major role in this cross-sectional analysis. In our study, smoking but not alcoholism correlated with severity of DR. Hammes et al study revealed significant correlation of glycaemic control, HDL-cholesterol and diastolic blood pressure on the occurrence of retinopathy. An Italian study showed Systolic BP was significantly related to Diabetic retinopathy, but Diastolic BP was not significantly related, and our study showed similar results.

#### Conclusion

Diabetic retinopathy is a significant complication of diabetes mellitus, which is a cause of preventable blindness or Sight Threatening Retinopathy (STDR). Prevalence of diabetic retinopathy among diabetic patients in our study is 31.81%. Average age of Diabetic retinopathy patients is around 54 years. Several studies

correlated development of DR and severity with duration of DM, systolic and diastolic hypertension, BMI, albuminuria, smoking, alcoholism and decreased vitamin D levels. Our study correlated with statistically significant figures with regards to Systolic blood pressure, albuminuria and smoking. Other factors like female preponderance, higher BMI, diastolic blood pressure and alcohol intake were high among patients of diabetic retinopathy, but the values did not show statistically significant correlation with the severity of diabetic retinopathy.

#### References

1. Ghafour IM, Allan, Foulds WS. Common causes of blindness and visual handicap in the west of Scotland. Br. J Ophthalmol 1983; 67 (4) :209-13.
2. Sjolie AK. Ocular complications in insulin treated diabetes mellitus. Ophthalmology 1985; 63:2-76.
3. Kahn HA, Hiller R. Blindness caused by diabetic retinopathy. Am J Ophthalmol. 1974;87 (1): 58-67.
4. Dandona L, Dandona R, Naduvilah TJ, McCarty CA, Rao GN. Population based assessment of diabetic retinopathy in an urban population in southern India. Br Ophthalmol 1999; 83: 936-40.
5. Bender AP, Sprafka JK, Jagger HG. A population based case control study of diabetic complications in Wadena. Marshal and Grand Rapids, Minnesota eds. Diabetes 1984; 73: 92A.
6. Mohan V, Vijayprabha R, Rena M. Vascular complications in long term South Indian NIDDM of over 25 years duration. Diabetes and Clin Pract 1996;31: 133-40.
7. Klein BEK, Klein R, Moss SE. Is increased blood pressure associated with Diabetic retinopathy? Diabetes 1984; 33: 45A.
8. United Kingdom Prospective Diabetes Study (UKPDS) Group. High Blood Pressure Control and risk of micro vascular and macro vascular complication in type 2 diabetes. UKPDS 38. BMJ, 1998; 317:703-13.
9. Dorman TL, Carter RD, Bron AJ, Turner RC, Mann JI. Low density lipoprotein cholesterol: An association with the severity of diabetic retinopathy. Diabetologia 1982; 22: 167-70.
10. Ferris FL III, Chew KY, Hoogwerf BJ. Serum lipids and diabetic retinopathy. Early Treatment Diabetic Retinopathy Study Research Group. Diabetes Care 1996; 19: 1291-93.

**Conflict of Interest: Nil**

**Source of support: Nil**